

Normable and metrizable locally convex spaces

(EXERCISES FOR LECTURE 3)

- 3.1.** Let S be an infinite set. Show that there are no continuous norms on \mathbb{K}^S . As a corollary, \mathbb{K}^S is not normable.
- 3.2.** Let X be a noncompact, completely regular (i.e., Tychonoff) topological space. Show that there are no continuous norms on $C(X)$. As a corollary, $C(X)$ is not normable.
- 3.3.** Let $U \subset \mathbb{R}^n$ be a nonempty open set. Show that there are no continuous norms on $C^\infty(U)$. As a corollary, $C^\infty(U)$ is not normable.
- 3.4.** Let $U \subset \mathbb{C}^n$ be a nonempty open set. When does $\mathcal{O}(U)$ have a continuous norm?
- 3.5.** Show that the following spaces are not normable, although each of them has a continuous norm:
(a) s ; (b) $C^\infty[a, b]$; (c) $\mathcal{S}(\mathbb{R}^n)$; (d) $\mathcal{O}(U)$ (where U is a nonempty domain in \mathbb{C}).
- 3.6.** Prove that the following spaces are metrizable:
(a) $C(X)$, where X is a second countable, locally compact topological space;
(b) all spaces from Exercises 3.3–3.5.
- 3.7.** Let S be a set. Show that \mathbb{K}^S is metrizable if and only if S is at most countable.
- 3.8.** Show that a vector space equipped with the strongest locally convex topology is metrizable if and only if it is finite-dimensional.
- 3.9.** Let X be a normed space. Show that
(a) the dual space X' equipped with the weak* topology is metrizable if and only if the dimension of X is at most countable;
(b) X equipped with the weak topology is metrizable if and only if it is finite-dimensional.
- 3.10*.** Let X be a finite-dimensional vector space. Show that there is only one topology on X which makes X into a Hausdorff topological vector space, and that this topology is determined by any norm on X . (This result was proved at the lectures in the special case of locally convex topologies.)
- 3.11*.** Prove that a Hausdorff topological vector space is finite-dimensional if and only if it is locally compact.
- 3.12*.** Prove that a topological vector space is semimetrizable if and only if its topology is generated by an F -seminorm. (This result was proved at the lectures in the special case of locally convex spaces.)